

MAVYRET
glecaprevir/pibrentasvir
100 mg/40 mg tablets

CRUSH, CONQUER, CURE

HEPATITIS C
IN ONLY 8
WEEKS.

For those new to treatment.

Cure means no hep C virus found in the blood 3 months after treatment ends. Individual results may vary.

USE

MAVYRET is a prescription medicine used to treat adults and children 3 years of age and older with chronic (lasting a long time) hepatitis C virus (hep C) genotypes 1, 2, 3, 4, 5 or 6 infection without cirrhosis or with compensated cirrhosis.

SAFETY CONSIDERATIONS

Do not take MAVYRET if you have certain liver problems or if you are taking the medicines atazanavir or rifampin.

Please see additional Important Safety Information throughout, including Boxed Warning on Hepatitis B reactivation.

Please see full Prescribing Information, including the Patient Information.

WHAT IS CHRONIC HEP C?

Hep C is a virus that multiplies daily and can damage the liver over time.

HEP C...



may not cause symptoms



can lead to liver damage

SAFETY CONSIDERATIONS (cont'd)

Hepatitis B virus (hep B) reactivation: Before starting treatment with MAVYRET, your doctor will do blood tests to check for hep B infection. If you have ever had hep B infection, hep B could become active again during or after treatment for hep C with MAVYRET.

WHAT IS MAVYRET?

MAVYRET stops the hep C virus from multiplying and CURES it.

MAVYRET...



can treat hep C in only 8 weeks for those new to treatment



is 3 pills taken at one time each day with food*

*For adults, adolescents 12 years and older, or children weighing at least 99 pounds (45 kg).

SAFETY CONSIDERATIONS (cont'd)

Hep B that becomes active again (called reactivation) may cause serious liver problems, including liver failure and death. Your doctor will monitor you if you are at risk for hep B reactivation during treatment and after you stop taking MAVYRET.

READY TO CURE HEP C?

Talk to your healthcare professional to learn more about MAVYRET.

MAVYRET CAN CURE MOST PEOPLE WITH HEP C AND...



can fit into your daily routine



can be taken with medication-assisted treatment for opioid use disorder* and most birth control methods

Your healthcare professional can tell you if it is safe to take MAVYRET with your birth control.

*If you are taking naltrexone, talk to your healthcare professional to learn if you can still take MAVYRET.

WHAT DOES MAVYRET COST?

Cost shouldn't get in the way of getting treated.

MAVYRET IS COVERED BY MOST INSURANCE...



Medicaid: pay \$20 or less per month for MAVYRET, depending on state plan



Commercial insurance: may pay as little as \$5 per month with MAVYRET savings card

myAbbvie Assist

myAbbvie Assist: if you don't have health insurance, AbbVie may be able to help



TO FIND OUT WHAT MAVYRET
MAY COST FOR YOU, CALL
1-877-628-9738.

Ask your healthcare professional about MAVYRET and CONQUER hep C.

Learn how to
CRUSH
hep C here:



SAFETY CONSIDERATIONS (cont'd)

What should I tell my doctor before taking MAVYRET?

If you have had hep B infection, have liver problems other than hep C infection, have HIV-1 infection, have had a liver or a kidney transplant, and all other medical conditions.

abbvie

MAVYRET
glecaprevir/pibrentasvir
100 mg/40 mg tablets

© 2023 AbbVie. All rights reserved.
MAVYRET® and its design are registered trademarks of AbbVie Inc.

US-MAVY-230153

“ I once said I
HAVE hep C.
Now I can say
I HAD hep C. ”

—Danny, CURED
with MAVYRET

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about MAVYRET?

Hepatitis B virus (hep B) reactivation: Before starting treatment with MAVYRET, your doctor will do blood tests to check for hep B infection. If you have ever had hep B infection, hep B could become active again during or after treatment for hep C with MAVYRET. Hep B that becomes active again (called reactivation) may cause serious liver problems, including liver failure and death. Your doctor will monitor you if you are at risk for hep B reactivation during treatment and after you stop taking MAVYRET.

Do not take MAVYRET if you:

- Have certain liver problems
- Are taking the medicines atazanavir or rifampin

What should I tell my doctor before taking MAVYRET?

- If you have had hep B infection, have liver problems other than hep C infection, have HIV-1 infection, have had a liver or a kidney transplant, and all other medical conditions.
- If you are pregnant or plan to become pregnant, or if you are breastfeeding or plan to breastfeed. It is not known if MAVYRET will harm your unborn baby or pass into your breast milk. Talk to your doctor about the best way to feed your baby if you take MAVYRET.

- **About all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. MAVYRET and other medicines may affect each other. This can cause you to have too much or not enough MAVYRET or other medicines in your body. This may affect the way MAVYRET or your other medicines work or may cause side effects.

- **Do not start taking a new medicine without telling your doctor.** Your doctor can tell you if it is safe to take MAVYRET with other medicines.

What are the possible side effects of MAVYRET?

- **In people who had or have advanced liver problems before starting treatment with MAVYRET, there is a rare risk of worsening liver problems, liver failure, and death.** Your doctor will check you for signs and symptoms of worsening liver problems during treatment with MAVYRET. Tell your doctor right away if you have any of the following: nausea; tiredness; yellowing of your skin or white part of your eyes; bleeding or bruising more easily than normal; confusion; dark, black, or bloody stool; loss of appetite; diarrhea; dark or brown (tea-colored) urine; swelling or pain on the upper right side of your stomach area (abdomen); sleepiness; vomiting of blood; or lightheadedness.

- The most common side effects of MAVYRET are headache and tiredness.

These are not all the possible side effects of MAVYRET. Call your doctor for medical advice about side effects.

This is the most important information to know about MAVYRET. For more information, talk to your doctor or healthcare provider.

MAVYRET oral pellets are dispensed in unit-dose packets. Each packet contains 50 mg glecaprevir/20 mg pibrentasvir.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see full **Prescribing Information**, including the **Patient Information**.

If you are having difficulty paying for your medicine, AbbVie may be able to help. Visit AbbVie.com/myAbbVieAssist to learn more.

“ MAVYRET fit
into my daily
routine. ”

—Dianna, CURED
with MAVYRET

MAVYRET is 3 pills taken at one time each day with food. Cure means no hep C virus found in the blood 3 months after treatment ends. Individual results may vary.

#1 PRESCRIBED MEDICINE
FOR CHRONIC HEP C*

*Based on data from IQVIA, week ending
1/12/2018 to 8/11/2023.



Ready to hear
more from other
real patients?

Visit MAVYRETstories.com
to learn how others have
CONQUERED and **CURED**
their hep C with MAVYRET.

MAVYRET
glecaprevir/pibrentasvir
100 mg/40 mg tablets

Cure means no hep C virus found in the blood 3 months after treatment ends. Individual results may vary.