



CRUSH
YOUR CHRONIC
HEP C IN ONLY
8 WEEKS*

*For those new to treatment.

USE

MAVYRET is a prescription medicine used to treat adults and children 3 years of age and older with chronic (lasting a long time) hepatitis C virus (hep C) genotypes 1, 2, 3, 4, 5 or 6 infection without cirrhosis or with compensated cirrhosis.

SAFETY CONSIDERATIONS

Do not take MAVYRET if you have certain liver problems or if you are taking the medicines atazanavir or rifampin

Please see full [Prescribing Information](#), including the [Patient Information](#).

WHY CURE[†]
your hep C now?



The hep C virus is busy multiplying a trillion times a day and has the potential to do serious damage to your liver



This may lead to serious health issues, including:

- Cirrhosis
- The need for a liver transplant
- Liver cancer

[†]Cure means no hep C virus found in the blood 3 months after treatment ends. Individual results may vary.

SAFETY CONSIDERATIONS (cont'd)

Hepatitis B virus (hep B) reactivation: Before starting treatment with MAVYRET, your doctor will do blood tests to check for hep B infection. If you have ever had hep B infection, hep B could become active again during or after treatment for hep C with MAVYRET.

WHY GET CURED
with MAVYRET?



MAVYRET stops the hep C virus and CURES it

98%

MAVYRET has a 98%[‡] CURE rate

8
WEEKS

It only takes 8 weeks[§]



For adults: It's only taken once a day—3 pills with food

[‡]Studies showed cure rates varied from 95% to 99% in 8 weeks for those new to treatment.

[§]For those new to treatment.

SAFETY CONSIDERATIONS (cont'd)

Hep B that becomes active again (called reactivation) may cause serious liver problems, including liver failure and death. Your doctor will monitor you if you are at risk for hep B reactivation during treatment and after you stop taking MAVYRET.

What to know about
TAKING MAVYRET



MAVYRET can be taken with **medication-assisted treatment for opioid-use disorder**^{||}




MAVYRET can be taken with **many types of birth control**^{||}



Some people have side effects—the most common are headache and tiredness

^{||} If you are taking naltrexone, talk to your healthcare professional to learn if you can still take MAVYRET.

[†] Your healthcare professional can tell you if it is safe to take MAVYRET with your birth control.



Cost shouldn't get in
THE WAY OF CURE



MAVYRET is covered by most types of insurance



Most people with Medicaid pay \$5 or less for MAVYRET, depending on their state plan



People with commercial insurance pay as little as \$5 a month with the MAVYRET Savings Card



If you don't have insurance, myAbbVie Assist may be able to help. Call 1-800-222-6885



WITH MAVYRET, YOU'RE NOT ALONE—CALL 1-877-628-9738

A Nurse Ambassador is there to answer questions and help navigate the insurance process.**

**Ambassadors are provided by AbbVie and do not provide medical advice or work under the direction of the prescribing healthcare professional (HCP). They are trained to direct patients to speak with their HCP about any treatment-related questions, including further referrals.

ASK YOUR DOCTOR
for MAVYRET



Get
CURED.
It's an incredible feeling. ■■

—Dianna, CURED with MAVYRET

#1 PRESCRIBED MEDICINE FOR CHRONIC HEPATITIS C^{††}

^{††}Based on data from IQVIA, week ending 1/12/2018 to 8/11/2023.

SAFETY CONSIDERATIONS (cont'd)

What should I tell my doctor before taking MAVYRET? If you have had hep B infection, have liver problems other than hep C infection, have HIV-1 infection, have had a liver or a kidney transplant, and all other medical conditions.



IMPORTANT SAFETY INFORMATION AND USE

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about MAVYRET?

Hepatitis B virus (hep B) reactivation: Before starting treatment with MAVYRET, your doctor will do blood tests to check for hep B infection. If you have ever had hep B infection, hep B could become active again during or after treatment for hep C with MAVYRET. Hep B that becomes active again (called reactivation) may cause serious liver problems, including liver failure and death. Your doctor will monitor you if you are at risk for hep B reactivation during treatment and after you stop taking MAVYRET.

Do not take MAVYRET if you:

- Have certain liver problems
- Are taking the medicines atazanavir or rifampin

What should I tell my doctor before taking MAVYRET?

- If you have had hep B infection, have liver problems other than hep C infection, have HIV-1 infection, have had a liver or a kidney transplant, and all other medical conditions.
- If you are pregnant or plan to become pregnant, or if you are breastfeeding or plan to breastfeed. It is not known if MAVYRET will harm your unborn baby or pass into your breast milk. Talk to your doctor about the best way to feed your baby if you take MAVYRET.

- **About all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. MAVYRET and other medicines may affect each other. This can cause you to have too much or not enough MAVYRET or other medicines in your body. This may affect the way MAVYRET or your other medicines work or may cause side effects.

- **Do not start taking a new medicine without telling your doctor.** Your doctor can tell you if it is safe to take MAVYRET with other medicines.

What are the possible side effects of MAVYRET?

- **In people who had or have advanced liver problems before starting treatment with MAVYRET, there is a rare risk of worsening liver problems, liver failure, and death.** Your doctor will check you for signs and symptoms of worsening liver problems during treatment with MAVYRET. Tell your doctor right away if you have any of the following: nausea; tiredness; yellowing of your skin or white part of your eyes; bleeding or bruising more easily than normal; confusion; dark, black, or bloody stool; loss of appetite; diarrhea; dark or brown (tea-colored) urine; swelling or pain on the upper right side of your stomach area (abdomen); sleepiness; vomiting of blood; or lightheadedness.

- The most common side effects of MAVYRET are headache and tiredness.

These are not all the possible side effects of MAVYRET. Call your doctor for medical advice about side effects.

This is the most important information to know about MAVYRET. For more information, talk to your doctor or healthcare provider.

MAVYRET oral pellets are dispensed in unit-dose packets. Each packet contains 50 mg glecaprevir/20 mg pibrentasvir.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see full [Prescribing Information](#), including the [Patient Information](#).


If you are having difficulty paying for your medicine, AbbVie may be able to help. Visit AbbVie.com/myAbbVieAssist to learn more.

USE

MAVYRET is a prescription medicine used to treat adults and children 3 years of age and older with chronic (lasting a long time) hepatitis C virus (hep C):

- Genotypes (GT) 1, 2, 3, 4, 5 or 6 infection without cirrhosis or with compensated cirrhosis
- Or GT 1 infection and have been previously treated with a regimen that contained a hep C NS5A inhibitor or an NS3/4A protease inhibitor, but not both.

WHAT DO REAL MAVYRET patients say?



“Taking MAVYRET once a day was **REALLY EASY FOR ME.**”

—Rick

“I GOT CURED WHILE IN RECOVERY.”

MAVYRET was my way to move on from hep C.

—Dianna



“I used heroin for 10 years. Curing my hep C helped me **MOVE BEYOND MY PAST.**”

—James



“Getting CURED with MAVYRET was **A BURDEN OFF MY SHOULDERS.**”

—Jeanne



HEAR MORE FROM real MAVYRET patients

MAVYRET
glecaprevir/pibrentasvir
100 mg/40 mg tablets

“**YOU GOTTA GET CURED — there's no reason to live with hep C.**”

—Jules, CURED with MAVYRET